

TABLE 2
ALCOHOL-INDUCED DEATHS
CAUSE-SPECIFIC DEATH RATES
CALIFORNIA, 2002
(By Place of Residence)

CAUSE OF DEATH	ICD-10 CODES	2002 DEATHS	CRUDE RATE	AGE-ADJUSTED RATE	95% CONFIDENCE LIMITS	
					LOWER	UPPER
Mental and behavioural disorders due to alcohol use	F10	844	2.357	2.470	2.303	2.637
Degeneration of nervous system due to alcohol	G31.2	7	0.020 *	0.022 *	0.006	0.038
Alcoholic polyneuropathy	G62.1	1	0.003 *	0.003 *	0.000	0.008
Alcoholic cardiomyopathy	I42.6	90	0.251	0.264	0.209	0.319
Alcoholic gastritis	K29.2	4	0.011 *	0.012 *	0.000	0.023
Alcoholic fatty liver	K70.0	39	0.109	0.111	0.076	0.145
Alcoholic hepatitis	K70.1	122	0.341	0.351	0.289	0.414
Alcoholic cirrhosis of liver	K70.3	1,663	4.645	4.922	4.684	5.159
Alcoholic hepatic failure	K70.4	234	0.654	0.681	0.594	0.769
Alcoholic liver disease, unspecified	K70.9	437	1.221	1.278	1.158	1.398
Accidental poisoning by and exposure to alcohol	X45	46	0.128	0.130	0.093	0.168
Intentional self poisoning by and exposure to alcohol	X65	7	0.020 *	0.020 *	0.005	0.035
Poisoning by and exposure to alcohol, undetermined intent	Y15	3	0.008 *	0.006 *	0.000	0.013

Note : Cause-specific death rates are per 100,000 population (n=35,802,238).

Alcoholic fibrosis and sclerosis of liver (K70.2) and finding of alcohol in blood (R78.0) were omitted from this table because of zero events.

* Death rate unreliable, relative standard error is greater than or equal to 23 percent.

Source : State of California, Department of Finance; 2002 Population: Population Projections by Age, Race/Ethnicity and Sex, December 1998.
State of California, Department of Health Services, Death Records.